inSite Digestive Healthcare COLONOSCOPY PREPARATION SHEET: GENERIC MODEL

Facility:	□ Check with	your gastro	office if you're	not CERTAIN	WRITE IT DOWN HERE:

Procedure Date:	Check In Time:

5 Days Prior	4 Days Prior	3 Days Prior	2 Days Prior	One Day Prior	Colonoscopy Day
ride to and from Facility Go to pharmacy with your prescription or list of laxatives and pack of bisacodyl 5mg tablets For sore bottom:plain or aloe baby wipesDesitin or A&D ointment Stop taking Coumadin or Plavix. Follow the instructions of		□ Review diet instructions for the coming 3 days. Plan meals and shop to comply with instructions □ Confirm your ride to and from facility □ Stop any fiber supplements □ Eat well balanced meals	clear liquid during the day Eat well balanced meals BUT AVOID: Raw fruits Raw vegetables Nuts or popcorn Foods with small seeds such as tomato, sesame or melon seeds No solid food after midnight Ies Orn Stop taking Eliquis, Pradaxa, Xarelto, Savasa	□ Drink at least 8 glasses of water or clear liquid during the day □ Follow a strict clear diet ALL DAY □ Take 1 Dulcolax (bisacodyl) 5 mg tablet at 12 noon. Prepare laxative drinks you'll need, set aside □ Starting at 5-6pm, drink prep fluids as directed & take one more Dulcolax (bisacodyl) 5 mg	□ You may take regular prescriptions with small sips of water (but no blood thinners) □ If you take insulin shots, follow the instructions for lowered dose
	- Raw fru - Raw ve - Nuts or - Foods v small se	BUT AVOID: - Raw fruits - Raw vegetables - Nuts or popcorn - Foods with small seeds such as tomato, sesame		tablet ☐ If no bowel movement occurs in 4 hours, continue to drink extra prep fluid ☐ Verify that you have info you'll need to bring with you to facility	 □ Be sure to bring: Insurance card or Medicare card Identification (CA driver's license, or ID card) Cash, check or credit card (Visa
alternative blood thinners are needed. Do not proceed without	or Plavix for 5 days	or melon seeds	dose (48 hours prior to procedure).	□ Do not schedule any important activities on the day of procedure	
talking to your Gastro doctor if taking Brilinta or Effient			□Do NOT stop low dose (81mg) aspirin taken for heart or stroke reasons	□ You may not drive or go home unaccompanied in a taxi or UBER etc after your procedure	□ Allow 2-3 hours at the facility before your driver will pick you up. Staff will call your driver to coordinate
		•	nded) Aspirin or NSAID kay or meds for migra	•	□Normal diet after you are home

inSite Digestive Healthcare COLONOSCOPY PREPARATION SHEET: GENERIC MODEL

Facility: Check with your gastro office if you're not CERTAIN WRITE IT DOWN HERE:						
Procedure Date:		Check In Time:_				

Two days prior: Consider stopping all casual (not Dr. recommended) Aspirin or NSAIDS-no ibuprofen, Advil, Aleve, Naprosyn, Excedrin---only **Tylenol is okay or meds for migraine**